

Depression & Anxiety

Treatment approaches covered

Psychopharmacotherapy | Psychotherapy | Lifestyle interventions | Self-guided help | Other forms of therapy



Details

- 1 Recommended to track via the Patient Health Questionnaire (PHQ-9)
- 2 Includes symptoms of general anxiety, social phobia, agoraphobia, post-traumatic stress disorder, panic disorder, and obsessive-compulsive disorder
- 3 Recommended to track via the Generalized Anxiety Disorder (GAD-7), and for those with specific anxiety disorders: the Social Phobia Inventory (SPIN), the Mobility Inventory for Agoraphobia (MIA), the Impact of Event Scale - Revised for Post-Traumatic Stress Disorder (IES-R), the Panic Disorder Severity Scale (PDSS-SR), and the Obsessive-Compulsive Inventory (OCI-R)
- 4 Recommended to track via the World Health Organization Disability Assessment 2.0 (WHODAS 2.0)
- 5 Includes work status and disease-related absenteeism

For a complete overview of this Set, including definitions for each measure, time points for collection, and associated risk factors, visit <https://connect.ichom.org/patient-centered-outcome-measures/depression-anxiety/>


Contributors

For more information about the process of developing a Set of Patient-Centered Outcome Measures, visit ichom.org/how-we-work/

The Sponsors



The Working Group



Australia

Maria Kangas | Centre for Emotional Health, Macquarie University

Brazil

Roberta Alvarenga Reis | Universidade Federal do Rio Grande do Sul

Canada

Anne Crocker | Douglas Mental Health University Institute, McGill University
Lucie Langford*
Alain Lesage | University of Montreal

Chile

Graciela Rojas | University of Chile

Germany

Sandra Nolte | Charité Universitätsmedizin Berlin
Alexander Obbarius | Charité Universitätsmedizin Berlin
Matthias Rose | Charité Universitätsmedizin Berlin

India

Vikram Patel | Centre for Chronic Conditions and Injuries; Public Health Foundation of India

Japan

Toshi A. Furukawa | Kyoto University

Netherlands

Edwin de Beurs | Stichting Benchmark GGZ
Paul Emmelkamp | University of Amsterdam

Sweden

Erik Hedman | Karolinska Institutet

Uganda

Doris M. Mwesigire | Makerere University

United Kingdom

David M. Clark | University of Oxford
David Smithson* | Anxiety UK

United States

Lee Baer | Harvard Medical School; Massachusetts General Hospital
Kelly Woolaway-Bickel | Department of the U.S. Army, Office of the Surgeon General
Paul A. Pilkonis | University of Pittsburgh
Harold A. Pincus | Columbia University
Cathy Sherbourne | The RAND Corporation

*Patient representative